



# Size Distribution and CCN Activation Ratio of Bacteria Ghosts

Alexa Otto<sup>1</sup> ([alexa.otto@und.edu](mailto:alexa.otto@und.edu)), Johannes Kassmannhuber<sup>2</sup>, Werner Lubitz<sup>2</sup>, and David Delene<sup>1</sup>



<sup>1</sup>University of North Dakota, Grand Forks, North Dakota; <sup>2</sup>Bird-C, Vienna, Austria

## Overview

Bacteria ghosts (BG) are non-living bacterial cell envelopes of Gram-negative bacteria. BG carrying ice nucleation proteins (INP-BG) have been shown to be effective ice nuclei. Experiments are being conducted at the University of North Dakota to collect lab data such as the size distribution and cloud condensation nuclei (CCN) activation ratio. The CCN activation ratio is an indication that the bacteria ghosts could be a good immersion ice nuclei.






Image showing the instrument setup for obtaining the size distribution and CCN activation Size of bacteria ghosts (BG).

## Acknowledgments


The National Institute of General Medical Sciences of the National Institutes of Health under Award Number T34GM122835 supported Alexa Otto's work on this research. The content present here is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.



Microscopy images of the E. Coli C 41 bacteria ghosts.



Scanning Mobility Particle Sizer (SMPS) measurements of the E. Coli C41 bacteria ghosts.



The particle size spectrum of the E. Coli C41 bacteria ghosts using a Aerodynamic Particle Sizer (APS).




Image showing the CCN activation ratio of the E. Coli C41 bacteria ghosts, which is obtain from concurrent Cloud Condensation Nuclei (CCN) counter and Condensation Particle Counter (CPC) measurements. The mean CCN activation ratio is 0.44 +/- 0.03.

## Results

- The size distribution peaks at 70 nm and has another peak at 0.7 μm.
- Approximately 44 percent of the bacteria ghosts activates as cloud condensation nuclei which indicates the potential for good immersion nuclei.

## Future Work

- Create code to combine the APS and SMPS data to better combine the two size peaks.
- Compare E. Coli C41 bacteria ghosts and E. Coli C41 bacteria ghosts with the InaZ protein.
- Determine the activation ratio as a function of particle size of the bacteria ghosts.